

NAS SYMPTOMS (n=135)	Frequency (%)
COMMON	
Tremors	
- - Mild/disturbed	96
- - Mild/undisturbed	95
- - Marked/disturbed	77
- - Marked/undisturbed	67
High-pitched cry	95
Continuous high-pitched cry	54
Sneezing	83
Increased muscle tone	82
Frantic sucking of fists	79
Regurgitation	74
Sleeps less than 3 hr after feeding	65
Sleeps less than 2 hr after feeding	66
Sleeps less than 1 hr after feeding	58
Respiratory rate greater than 60/min	66
Poor feeding	65
Hyperactive Moro reflex	62
Loose stools	51
LESS COMMON	
Sweating	49
Excoriation	43
Mottling	33
Nasal stuffiness	33
Frequent yawning	30
Fever less than 101°F (39.3°C)	29
Respiratory rate greater than 60/min and retractions	28
Markedly hyperactive Moro reflex	15
Projectile vomiting	12
Watery stools	12
Fever less than 101°F (39.3°C)	3
Dehydration	1
Generalized convulsions	1

Finnegan L. *Neonatal abstinence syndrome: assessment and pharmacotherapy*. Neonatal Therapy: An update, F. F. Rubaltelli and B. Granti, editors. Elsevier Science Publishers B. V. (Biomedical Division). 1986: 122-146 (4).